

ON UNIFORM CONVERGENCE AND A
SERIES OF VARIABLE TERMS

A THESIS

SUBMITTED TO THE FACULTY OF ATLANTA UNIVERSITY
IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR
THE DEGREE OF MASTER OF SCIENCE

BY

JAMES EDWARD DAWSON

DEPARTMENT OF MATHEMATICS

ATLANTA, GEORGIA

JANUARY 1964

$R = vi$ $T = 17$

ACKNOWLEDGMENT

The writer wishes to express his sincere appreciation to his advisor, Dr. Lonnie Cross, for his untiring service and words of encouragement, which made the writing of this thesis possible.

PREFACE

This thesis is a brief study on the theory of uniform convergence. It contains material which ultimately leads to the analysis of a series of variable terms.

Chapter I contains basic information relative to uniform convergence needed in the remainder of the thesis. This information includes definitions and examples on uniform convergence.

Chapter II contains the analysis of a series of variable terms. The discussion of this series is the main objective of this thesis.

The information included in Chapter III gives a brief summary of the thesis, and it discusses the series considered in Chapter II. Furthermore, it discusses the series' connection with the theory of uniform convergence; that is also included in Chapter II.

TABLE OF CONTENTS

	Page
ACKNOWLEDGMENT.....	iii
PREFACE.....	iv
GLOSSARY OF SYMBOLS AND ABBREVIATIONS.....	vi
Chapter	
I. PRELIMINARY REMARKS ABOUT UNIFORM CONVERGENCE....	1
Uniform Convergence of Sequences.....	1
Definition.....	1
Negation of Uniform Convergence.....	2
Uniform Convergence of Series.....	3
Definition.....	3
II. SERIES OF VARIABLE TERMS.....	6
III. SUMMARY.....	13
BIBLIOGRAPHY.....	17

GLOSSARY OF SYMBOLS AND ABBREVIATIONS

\in "belongs to"

\Rightarrow "approaches"

\implies "implies"

\longrightarrow "tends to"

Q.E.D. (quod erat demonstrandum-"which was to be proved")

Σ "sigma," summation

CHAPTER I

PRELIMINARY REMARKS ABOUT UNIFORM CONVERGENCE

Uniform convergence of sequences.

Let $S_1(x), S_2(x), \dots, S_n(x), \dots$ be a sequence of functions defined on a set A . Each term of the sequence is a function of x and we denote it by $S_n(x)$. Thus $S_n(x)$ will converge on A if for every $x \in A$ the sequence of constants $\{S_n(x)\}$ converges. So then we assume $S_n(x)$ converges on A and define

$$S(x) \equiv \lim_{n \rightarrow +\infty} S_n(x).$$

Then the rapidity with which $S_n(x)$ approaches $S(x)$ can be expected to depend on the value of x .¹

Definition.—A sequence of functions $\{S_n(x)\}$, defined on a set A , converges uniformly on A to a function $S(x)$ defined on A , this being written

$$S_n(x) \rightrightarrows S(x),$$

if and only if corresponding to $\epsilon > 0$ there exists a number $N = N(\epsilon)$, dependent on ϵ alone and not on the point x , such that $n > N$ implies

$$|S_n(x) - S(x)| < \epsilon$$

for all $x \in A$.

¹J. M. H. Olmsted, Real Variables, New York, [1959], p. 270.

Example: Suppose $S_n(x) = \frac{x}{x+n}$. We want to show that

$S_n(x)$ converges uniformly on the interval $[0, b]$. So then for $\epsilon > 0$, we want to show there exists a number $N = N(\epsilon)$, such that

$$|S_n(x) - S(x)| < \epsilon$$

for all $n > N$ and $x \in [0, b]$. Observe that the

$$\lim_{n \rightarrow +\infty} S_n(x) = \lim_{n \rightarrow +\infty} \frac{x}{x+n} = 0.$$

Then we want $\left| \frac{x}{x+n} \right| < \epsilon$ for all $x \in [0, b]$ for $n > N$.

This implies that for $n > N$ $|x| < \epsilon |x+n| \Rightarrow \frac{|x|}{\epsilon} < |x+n| \Rightarrow$

$|x+n| > \frac{|x|}{\epsilon} \Rightarrow n > \frac{x}{\epsilon} (1-\epsilon)$. If $x = b$, then $n > \frac{b}{\epsilon} (1-\epsilon)$.

Take $N(\epsilon) \equiv \frac{b}{\epsilon} (1-\epsilon)$, and $\left| \frac{x}{x+n} \right| < \epsilon$ for all $n > N$ and $x \in [0, b]$.

Q.E.D.

Negation of uniform convergence.-A sequence of functions $\{S_n(x)\}$, defined on a set A , fails to converge uniformly on A to a function $S(x)$ defined on A if and only if there exists a positive number ϵ having the property that for any number N there exist a positive integer $n > N$ and a point $x \in A$ such that

$$|S_n(x) - S(x)| \geq \epsilon.$$

Example: Show that $S_n(x) = \frac{x}{x+n}$ does not converge uniformly on $[0, +\infty)$.

Solution: Suppose we take $\epsilon \equiv \frac{1}{2}$. If N is an arbitrary number, let us choose any $n > N$, and hold this n fixed. Next

we pick $x > n$. Then for this pair x and n ,

$$|S_n(x) - S(x)| = \frac{x}{x+n} > \frac{x}{x+x} = \frac{x}{2x} = \frac{1}{2}.$$

Since $\epsilon \equiv \frac{1}{2}$ this proves that $\frac{x}{x+n}$ does not converge uniformly on $[0, +\infty)$. Q.E.D.

Uniform convergence of series.

Let $U_1(x) + U_2(x) + \dots + U_n(x) + \dots$
be a series of functions defined on a set A , and let

$$S_n(x) \equiv U_1(x) + U_2(x) + \dots + U_n(x).$$

We say that this series of functions converges on A in case the sequence $\{S_n(x)\}$ converges on A . The series

$$U_1(x) + U_2(x) + \dots + U_n(x) + \dots$$

converges uniformly on A if and only if the sequence $\{S_n(x)\}$ converges uniformly on A .¹

Definition.—The series $\sum_{n=1}^{\infty} U_n(x)$ is uniformly convergent

to $S(x)$ for a set A of values of x if for each $\epsilon > 0$, an integer N can be found such that

$$|S_n(x) - S(x)| < \epsilon \text{ for } n \geq N \text{ and all } x \in A.^2$$

¹J. M. H. Olmsted, Real Variables, New York, [1959], p. 273.

²W. Kaplan, Advanced Calculus, Massachusetts, [1952], p. 339.

Thus we see that the uniform convergence of the series $\sum U_n(x)$ is equivalent to uniform convergence of its sequence of partial sums $S_n(x)$.

At this point, we will demonstrate the definition stated above with the use of two complex series $\sum_{n=0}^{\infty} z^n$ when $|z| < 1$ and $\sum_{n=0}^{\infty} \frac{1}{z^n}$ when $|z| > 1$.

This can be done since the interchange of interval for region in dealing with complex numbers does not destroy the essence of the definition.¹

These two series will be used later in this study to show that a series of variable terms converges to a function of z when $|z| < 1$ and also when $|z| > 1$.

Example 1. Show that the series $\sum_{n=0}^{\infty} z^n$ converges uniformly when $|z| < \rho$ for $\rho < 1$.

Consider $S_n(z) = \sum_{k=0}^n z^k = 1 + z + z^2 + z^3 + \dots + z^n = \frac{1}{1-z} - \frac{z^{n+1}}{1-z}$. Moreover, $\lim_{n \rightarrow +\infty} S_n(z) = S(z)$ when $|z| < 1$. Therefore, $\lim_{n \rightarrow +\infty} \frac{1}{1-z} - \frac{z^{n+1}}{1-z} = \frac{1}{1-z}$ when $|z| < 1$. By the definition

of uniform convergence, for $\epsilon > 0$, there must exist a positive

¹K. Knopp, Theory and Application of Infinite Series, New York, [1928], p. 428.

integer $N = N(\epsilon)$ such that $|S_n(z) - S(z)| < \epsilon$ when $n > N$ and all z such that $|z| \leq \rho$.

$$\text{Observe } |S_n(z) - S(z)| = \left| \frac{1}{1-z} - \frac{z^{n+1}}{1-z} - \frac{1}{1-z} \right| = \left| \frac{z^{n+1}}{1-z} \right| =$$

$$\left| \frac{z^{n+1}}{1-z} \right|. \text{ Take } N = N(\epsilon) \equiv \frac{\epsilon}{1-\rho}, \text{ and } \left| \frac{z^{n+1}}{1-z} \right| < \epsilon \text{ for all}$$

$n > N$ and all z such that $|z| \leq \rho$. Therefore, the $\sum_{n=0}^{\infty} z^n$ converges uniformly in $|z| \leq \rho$. Q.E.D.

Example 2. Show that the $\sum_{n=0}^{\infty} \frac{1}{z^n}$ converges uniformly in $|z| \geq \rho$ for ρ arbitrary and $\rho > 1$.

$$\text{Consider } S_n(z) = \sum_{k=0}^n \frac{1}{z^k} = 1 + \frac{1}{z} + \frac{1}{z^2} + \dots + \frac{1}{z^n} = \frac{z}{z-1} -$$

$$\frac{z-1}{z^n}. \text{ Thus } \lim_{n \rightarrow +\infty} \frac{z}{z-1} - \frac{z-1}{z^n} = \frac{z}{z-1} \text{ for all } z \text{ such that}$$

$$|z| > 1.$$

We want to show for $\epsilon > 0$, there exists a positive integer $N = N(\epsilon)$ such that for $n > N$ $|S_n(z) - S(z)| < \epsilon$.

$$\text{Observe } \left| \frac{z}{z-1} - \frac{z-1}{z^n} - \frac{z}{z-1} \right| = \left| -\frac{z-1}{z^n} \right| = \left| \frac{z-1}{z^n} \right|. \text{ If } |z| = \rho,$$

$$\text{then take } N(\epsilon) \equiv \frac{\epsilon}{\rho^n}. \text{ Now } \left| \frac{z-1}{z^n} \right| < \epsilon \text{ for all } n > N \text{ and all}$$

z such that $|z| \geq \rho$. Hence the $\sum_{n=0}^{\infty} \frac{1}{z^n}$ converges uniformly

in $|z| \geq \rho$. Q.E.D.

CHAPTER II

SERIES OF VARIABLE TERMS

This chapter will deal with a series consisting of variable terms. Thus the value of the series depends on the choice of a definite quantity or variable. We shall consider a series whose terms depend on the variable z , that is, a series of the

form $\sum_{n=0}^{\infty} f_n(z)$. This series defines for certain values of z a function f . In particular we consider the series

$$\sum_{n=1}^{\infty} \frac{z^{n-1}}{(1-z^n)(1-z^{n+1})} \quad .^1 \quad \text{We shall show that it is equal to}$$

$$\frac{1}{(1-z)^2} \quad \text{when } |z| < 1, \text{ and is equal to } \frac{1}{z(1-z)^2} \quad \text{when } |z| > 1.$$

First we show that the series $\sum_{n=1}^{\infty} \frac{z^{n-1}}{(1-z^n)(1-z^{n+1})} = \frac{1}{(1-z)^2}$ when $|z| < 1$.

Suppose

$$S_n(z) = \sum_{k=1}^n \frac{z^{k-1}}{(1-z^k)(1-z^{k+1})}$$

is the sum of the first n terms. So then

¹E. T. Whittaker, Course of Modern Analysis, Cambridge, 1935, p. 59.

$$(1) \quad S_n(z) = \sum_{k=1}^n \frac{z^{k-1}}{(1-z^k)(1-z^{k+1})} = \frac{1}{(1-z)(1-z^2)} +$$

$$\frac{z}{(1-z^2)(1-z^3)} + \frac{z^2}{(1-z^3)(1-z^4)} + \frac{z^3}{(1-z^4)(1-z^5)} +$$

$$\frac{z^4}{(1-z^5)(1-z^6)} + \dots + \frac{z^{n-1}}{(1-z^n)(1-z^{n+1})}.$$

It follows that

$$(2) \quad S_n(z) = \frac{1}{(1-z)^2} \left\{ \frac{1}{(1+z)} + \frac{z}{(1+z)(1+z+z^2)} + \right.$$

$$\frac{z^2}{(1+z+z^2)(1+z+z^2+z^3)} + \frac{z^3}{(1+z+z^2+z^3)(1+z+z^2+z^3+z^4)} +$$

$$\frac{z^4}{(1+z+z^2+z^3+z^4)(1+z+z^2+z^3+z^4+z^5)} + \dots +$$

$$\left. \frac{z^{n-1}}{\left(\sum_{k=0}^{n-1} z^k\right) \left(\sum_{k=0}^n z^k\right)} \right\}.$$

Now it follows from algebra using the theorem on partial fractions, that the fraction $\frac{z}{(1+z)(1+z+z^2)} = \frac{A}{1+z} + \frac{Bz+C}{1+z+z^2}.$

Hence

$$z = A(1+z+z^2) + (Bz+C)(1+z)$$

$$= A + Az + Az^2 + Bz + Bz^2 + Cz + C$$

$$= (A+B)z^2 + (A+B+C)z + A+C.$$

Equating coefficients, we have

$$A + B = 0,$$

$$A + B + C = 1,$$

$$A + C = 0.$$

Hence $C = 1$, $A = -1$, and $B = 1$. Therefore,

$$\frac{z}{(1+z)(1+z+z^2)} = -\frac{1}{1+z} + \frac{z+1}{1+z+z^2}.$$

Consider now the fraction

$$\frac{z^2}{(1+z+z^2)(1+z+z^2+z^3)} = \frac{Az + B}{1+z+z^2} + \frac{Cz^2 + Dz + E}{1+z+z^2+z^3}.$$

Hence

$$\begin{aligned} z^2 &= (Az+B)(1+z+z^2+z^3) + (Cz^2+Dz+E)(1+z+z^2) \\ &= Az^4 + (A+B)z^3 + (A+B)z^2 + (A+B)z + B + Cz^4 + (C+D)z^3 + \\ &\quad (C+D+E)z^2 + (D+E)z + E \\ &= (A+C)z^4 + (A+B+C+D)z^3 + (A+B+C+D+E)z^2 + (A+B+D+E)z + (B+E). \end{aligned}$$

Equating coefficients, we have

$$A + C = 0,$$

$$A + B + C + D = 0,$$

$$A + B + D + E = 0,$$

$$A + B + C + D + E = 1,$$

$$B + E = 0.$$

Hence $A = -1$, $B = -1$, $C = 1$, $D = 1$, and $E = 1$. Therefore,

$$\frac{z^2}{(1+z+z^2)(1+z+z^2+z^3)} = -\frac{z+1}{1+z+z^2} + \frac{z^2+z+1}{1+z+z^2+z^3}.$$

Thus, it is easily seen that

$$\frac{z^{n-1}}{\left(\sum_{k=0}^{n-1} z^k\right) \left(\sum_{k=0}^n z^k\right)} = - \frac{\sum_{k=0}^{n-2} z^k}{\sum_{k=0}^{n-1} z^k} + \frac{\sum_{k=0}^{n-1} z^k}{\sum_{k=0}^n z^k}.$$

Substituting these results in equation (2), we obtain

$$(3) \quad S_n(z) = \frac{1}{(1-z)^2} \left\{ \frac{1}{1+z} - \frac{1}{1+z} + \frac{z+1}{1+z+z^2} - \right. \\ \left. \frac{z+1}{1+z+z^2} + \frac{z^2+z+1}{1+z+z^2+z^3} + \dots - \frac{\sum_{k=0}^{n-2} z^k}{\sum_{k=0}^{n-1} z^k} + \frac{\sum_{k=0}^{n-1} z^k}{\sum_{k=0}^n z^k} \right\}.$$

Since the first $2n-2$ terms of the sum (3) cancel each other leaving the last term, we have

$$S_n(z) = \frac{1}{(1-z)^2} \left\{ \frac{\sum_{k=0}^{n-1} z^k}{\sum_{k=0}^n z^k} \right\}.$$

Consider the sums $\sum_{k=0}^{n-1} z^k$ and $\sum_{k=0}^n z^k$ for $|z| < 1$.

Both converge to the same sum as $n \rightarrow +\infty$, namely $\frac{1}{1-z}$.

Hence, the

$$\lim_{n \rightarrow +\infty} \frac{1}{(1-z)^2} \left\{ \frac{\sum_{k=0}^{n-1} z^k}{\sum_{k=0}^n z^k} \right\} = \frac{1}{(1-z)^2},$$

when $|z| < 1$, for the expression $\left\{ \frac{\sum_{k=0}^{n-1} z^k}{\sum_{k=0}^n z^k} \right\} \rightarrow 1$ as $n \rightarrow +\infty$,

when $|z| < 1$. Q.E.D.

We will now show that the series

$$\sum_{n=1}^{\infty} \frac{z^{n-1}}{(1-z^n)(1-z^{n+1})} = \frac{1}{z(1-z)^2}$$

when $|z| > 1$. Consider the sum $\sum_{k=0}^n z^k$. Thus writing this

as $1 + z + z^2 + z^3 + \dots + z^n = \sum_{k=0}^n z^k$. Dividing this sum

through by z^n and multiplying through by z^n , we have

$$z^n \left(1 + \frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} + \dots + \frac{1}{z^n} \right) = \sum_{k=0}^n z^k.$$

Thus

$$\begin{aligned} z^n \sum_{k=0}^n \frac{1}{z^k} &= z^n \left(1 + \frac{1}{z} + \frac{1}{z^2} + \frac{1}{z^3} + \dots + \frac{1}{z^n} \right) \\ &= (1 + z + z^2 + \dots + z^n). \end{aligned}$$

But the series $\sum_{k=0}^{\infty} \frac{1}{z^k}$ converges for all z when $|z| > 1$.

Now observe

$$S_n(z) = \frac{1}{(1-z)^2} \left\{ \frac{z^{n-1} \sum_{k=0}^{n-1} \frac{1}{z^k}}{z^n \sum_{k=0}^n \frac{1}{z^k}} \right\} = \frac{1}{z(1-z)^2} \left\{ \frac{\sum_{k=0}^{n-1} \frac{1}{z z^k}}{\sum_{k=0}^n \frac{1}{z^k}} \right\}.$$

Since $\sum_{k=0}^{n-1} \frac{1}{z^k}$ and $\sum_{k=0}^n \frac{1}{z^k}$ both converge to the same limit

as $n \rightarrow +\infty$ when $|z| > 1$, namely $\frac{z}{z-1}$.

Therefore, it follows that

$$\lim_{n \rightarrow +\infty} \frac{1}{z(1-z)^2} \left\{ \frac{\sum_{k=0}^{n-1} \frac{1}{z^k}}{\sum_{k=0}^n \frac{1}{z^k}} \right\} = \frac{1}{z(1-z)^2}$$

when $|z| > 1$, for $\frac{\sum_{k=0}^{n-1} \frac{1}{z^k}}{\sum_{k=0}^n \frac{1}{z^k}} \rightarrow 1$ when $|z| > 1$ and $n \rightarrow +\infty$.

Hence,

$$\sum_{n=1}^{\infty} \frac{z^{n-1}}{(1-z^n)(1-z^{n+1})} = \frac{1}{z(1-z)^2} \text{ when } |z| > 1. \text{ Q.E.D.}$$

CHAPTER III

SUMMARY

We have seen from Chapter I the condition necessary for uniform convergence is that given $\epsilon > 0$, there exists a positive integer $N = N(\epsilon)$, such that the $|S_n(z) - S(z)| < \epsilon$ for all $n > N$ and z belongs to a region G . Thus we see that uniform convergence is defined on a set whereas convergence is defined at a point.

Now we will investigate what connection the series

$$\sum_{n=1}^{\infty} \frac{z^{n-1}}{(1-z^n)(1-z^{n+1})} \equiv \begin{cases} \frac{1}{(1-z)^2}, & \text{when } |z| < 1 \\ \frac{1}{z(1-z)^2}, & \text{when } |z| > 1 \end{cases}$$

have with the theory of uniform convergence.

It has been shown previously that this series converges for all z , such that $|z| < 1$ and $|z| > 1$. But outside of the regions $|z| < 1$ and $|z| > 1$, we have divergence of this series. Thus we see that the series

$$\sum_{n=1}^{\infty} \frac{z^{n-1}}{(1-z^n)(1-z^{n+1})}$$

converges to a definite sum, when z is restricted to $|z| < 1$ and $|z| > 1$.

Since

$$S_n(z) = \frac{1}{(1-z)^2} \left\{ \frac{\sum_{k=0}^{n-1} z^k}{\sum_{k=0}^n z^k} \right\},$$

convergence of this sequence depends totally on the behavior

of the sums $\sum_{k=0}^n z^k$ and $\sum_{k=0}^{n-1} z^k$. Since both behave similar, we

will consider one.

Observe

$$\lim_{n \rightarrow +\infty} \sum_{k=0}^n z^k = 1 + z + z^2 + \dots + z^n + \dots = \frac{1}{1-z},$$

when $|z| < 1$. This series converges for all z such that $|z| < 1$.

Since $\left\{ \sum_{k=0}^n z^k \right\}$ converges for all z such that the $|z| < 1$,

$$S_n(z) = \frac{1}{(1-z)^2} \left\{ \frac{\sum_{k=0}^{n-1} z^k}{\sum_{k=0}^n z^k} \right\}$$

converges for all z such that the $|z| < 1$. If z is bound

away from 1, then $S_n(z)$ will converge uniformly in this region; i.e., let $\rho < 1$ and arbitrary. Therefore, $S_n(z)$ converges uniformly for all z , such that $|z| \leq \rho$.¹

Since the sequence $\left\{ \sum_{k=0}^n z^k \right\}$ converges uniformly for all z

such that $|z| \leq \rho$, we have that $S_n(z)$ converges uniformly to

$$\frac{1}{(1-z)^2} \text{ when } |z| \leq \rho, \text{ for } \rho < 1.$$

Similarly, we have

$$S_n(z) = \frac{1}{z(1-z)^2} \left\{ \frac{\sum_{k=0}^{n-1} \frac{1}{z^k}}{\sum_{k=0}^n \frac{1}{z^k}} \right\}$$

converges for all z such that the $|z| > 1$. But convergence of this sequence depends wholly on the behavior of the sum

$\sum_{k=0}^n \frac{1}{z^k}$. Since this sum converges for all z such that the

$|z| > 1$, $S_n(z)$ converges for all z such that $|z| > 1$.

If $\rho > 1$ and if $|z| > \rho$, then $S_n(z)$ converges uniformly in $|z| > \rho$. For the sequence

¹K. Knopp, Theory of Function, New York, [1945], p. 72.

$\left\{ \sum_{k=0}^n \frac{1}{z^k} \right\}$ converges uniformly for all z such that the $|z| \geq \rho$

and $\rho > 1$. Hence $S_n(z)$ converges uniformly to $\frac{1}{z(1-z)^2}$ when

$|z| \geq \rho$. Q.E.D.

BIBLIOGRAPHY

John M. H. Olmsted, Real Variables, New York, Appleton-Century-Crofts, Inc., 1956.

W. Kaplan, Advanced Calculus, Massachusetts, 1952.

Konrad Knopp, Theory and Application of Infinite Series, London, Blackie and Son Limited, 1928.

E. T. Whittaker, A Course of Modern Analysis, Cambridge, At the University Press, 1935.

Konrad Knopp, Theory of Functions, New York, Dover Publications, 1945.